

**Case Report****Genomic Integration of HHV-6 Mimicking Viral Reactivation after Autologous Stem Cell Transplantation**

Rossana Mineri<sup>1</sup>, Jacopo Mariotti<sup>2</sup>, Barbara Sarina<sup>2</sup>, Lucio Morabito<sup>2</sup>, Roberto Crocchiolo<sup>2</sup>, Stefania Bramanti<sup>2</sup>, Tiziana Sarno<sup>1</sup>, Federica Tordato<sup>3</sup>, Carmelo Carlo-Stella<sup>2</sup>, Armando Santoro<sup>2,4</sup> and Luca Castagna<sup>2</sup>.

<sup>1</sup> Molecular Biology Section, Clinical Investigation Laboratory, Humanitas Clinical and Research Center, Via Manzoni 56, Rozzano 20089, Italy.

<sup>2</sup> Hematology Unit, Humanitas Cancer Center, Humanitas Clinical and Research Center, Via Manzoni 56, Rozzano 20089, Italy.

<sup>3</sup> Infectious Disease Unit, Humanitas Clinical and Research Center, Via Manzoni 56, Rozzano 20089, Italy.

<sup>4</sup> Humanitas University, Via Alessandro Manzoni 113, 20089 Rozzano (Mi), Italy.

**Competing interests:** The authors have declared that no competing interests exist.

**Abstract.** The monitoring of Human Herpesvirus 6 (HHV-6) after allogeneic stem cell transplantation has proven to be useful in preventing life-threatening complications; however, the pathogenic role of HHV-6 after autologous transplantation is not well-characterized, although viral reactivation might be responsible for significant complications even after this type of transplant. Here we report, for the first time to our knowledge, the case of a patient with chromosomally integrated HHV-6 (ciHHV-6), presenting with high titers of HHV-6 DNA copies after autologous transplantation, mimicking HHV-6 reactivation. The presence of viral DNA in the follicle bulb confirmed the ciHHV-6 and allowed for the discontinuation of the antiviral treatment. Due to the increasing awareness of HHV-6 potential pathogenicity and the fact that ciHHV-6 is expected in 1-2% of the population, such a case might be helpful in recognizing ciHHV-6, thus avoiding unnecessary and potentially toxic antiviral therapy once the viral genomic integration is confirmed.

**Keywords:** Bone marrow transplantation, Infectious disease, HHV-6 integration.

**Citation:** Mineri R., Mariotti J., Sarina B., Morabito L., Crocchiolo R., Bramanti S., Sarno T., Tordato F., Stella C.C., Santoro A., Castagna L. Genomic integration of HHV-6 mimicking viral reactivation after autologous stem cell transplantation. *Mediterr J Hematol Infect Dis* 2018, 10(1): e2018013, DOI: <http://dx.doi.org/10.4084/MJHID.2018.013>

**Published:** February 15, 2018

**Received:** October 3, 2017

**Accepted:** January 31, 2018

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Correspondence to: Rossana Mineri, PhD. Molecular Biology Section, Clinical Investigation Laboratory, Humanitas Clinical and Research Center, Via Manzoni 56, 20089 Rozzano, Italy. Tel. 0039 02 28844748. E-mail: [rossana.mineri@humanitas.it](mailto:rossana.mineri@humanitas.it)

**Introduction.** Human herpesvirus 6 (HHV-6) reactivation may be responsible for severe side effects after allogeneic hematopoietic stem cell transplantation (allo-HSCT) with early post-transplant mortality.<sup>1-3</sup> HHV-6 reactivation is typically detected 2 to 4 weeks after allo-HSCT and is associated with hepatitis, pneumonitis, CMV reactivation, fever, skin rash, myelosuppression, encephalitis, acute graft-versus-host disease; therefore, its regular

monitoring is recommended after those allo-HSCTs with a higher risk of reactivation, such as cord blood or haploidentical HSCT.<sup>4-5</sup> At our center, the control of HHV-6 is routinely performed after cord or haploidentical HSCT, whereas for all the other transplant settings (i.e., allo-HSCT from HLA-identical donor, or autologous stem cell transplantation, ASCT) the search for HHV-6 is made only in the presence of clinical conditions evoking a potential role of the


virus, such as prolonged aplasia or cutaneous rash<sup>6</sup>. Recently, it has been discovered that 1% to 2% of the population have chromosomally integrated HHV-6 (ciHHV-6), that is the presence of HHV-6 in every somatic cell leading to high copy numbers of the virus in the absence of any viral reactivation or disease.<sup>7-10</sup>

**Methods.** Here we present a case of a patient affected by non-Hodgkin's lymphoma undergoing high-dose chemotherapy followed by ASCT, with ciHHV-6 (detected through CMV HHV6,7,8 R-gene®, bioMérieux) mimicking viral reactivation early after transplant.

**Results.** A 66-years old man was diagnosed with stage IVA diffuse large B-cell lymphoma in December 2015 following the appearance of hepatomegaly, leading to the detection of a 7-cm hepatic mass by abdominal ultrasound. Complete staging showed sub- and supradiaphragmatic nodal and extranodal disease (liver, lung), with a Revised-International Prognostic Index (R-IPI) = 4 (stage, LDH higher than the average range, age > 60 years old and extranodal disease) and an NCCN IPI=5. The patient's history disclosed myocardial infarction 25 years before and concurrent HCV positivity. He received six cycles of full-dose R-CHOP with concomitant central nervous system prophylaxis, from April to July 2016. A new disease assessment in August 2016 showed metabolic complete remission. Due to the high-risk features at diagnosis, consolidation therapy with high-dose chemotherapy followed by ASCT

had been planned, and the patient was admitted to the transplantation unit at the end of August to receive ASCT, after conditioning chemotherapy with FEAM (fotemustine, etoposide, cytarabine, melphalan) regimen.<sup>11</sup> The aplastic phase was complicated by NCI-CTC grade 3 mucositis and Aspergillus pneumonia, classified as *probable*, according to EORTC criteria,<sup>12</sup> successfully treated with voriconazole (the patient has been receiving prophylaxis with fluconazole). On day +15 after ASCT, a determination of HHV-6 viral load was made because of persistent aplasia, showing  $9.7 \times 10^5$  copies/mL, with a total leucocytes count of  $1,380 \times 10^6/L$ . Molecular analysis by PCR was negative for CMV and EBV. Then, in the hypothesis of HHV-6 reactivation, known to have an incidence of 8-15% after ASCT,<sup>13-15</sup> antiviral therapy with foscarnet was started, and subsequent shift to ganciclovir was realized, after hematological recovery. However, the constant increase of HHV-6 load, observed in conjunction of the augmentation of the leucocyte counts (see **Figure 1**), led to the alternative hypothesis of ciHHV-6, soon confirmed by the analysis of patient's follicle bulb at the beginning of October.

Indeed, it is known that ciHHV-6 corresponds to the presence of one or more HHV-6 copies per white blood cell, with high HHV-6 titers persisting over time in blood, and of virus in tissues, allowing for a diagnosis made through the analysis of follicle bulb or nails.<sup>16</sup> Nonetheless, a confirmatory test is needed in the diagnostic process, as we did in the present case. The patient



**Figure 1.** Whole blood HHV-6 titers and total leucocyte counts are shown here. On the X axis, the days after autologous transplant are shown.

was discharged in good clinical conditions at day +28 from ASCT, and the antiviral treatment was discontinued at the moment of hospital discharge. Finally, the cause of the prolonged aplasia was not identified. Three months later the patient was healthy, and a CT scan confirmed the complete remission and a favorable evolution of the fungal infection. As expected, leukocyte, neutrophil and platelet counts were normalized, while HHV-6 blood titer was  $1.14 \times 10^7$  copies/mL with a total leucocyte count of  $6,920 \times 10^6/L$  in early January 2017.

**Discussion.** The present clinical case shows a genomic integration of HHV-6 mimicking reactivation early after autologous stem cell transplantation. Similar cases of transmission of integrated HHV-6 were described in recipients of allogeneic transplant from HLA matched or mismatched related donors;<sup>17-18</sup> more recently, it has been described after allogeneic, combined cord blood/haplo transplantation.<sup>19</sup> However, to our knowledge, our case is the first described after autologous stem cell transplantation. It is known from the literature that approximately 95% of the healthy adults experienced the HHV-6 infection in childhood. After primary infection, HHV-6 persists in the host and is detectable in multiple tissues, such as salivary glands, brain cells, monocytes, and early bone marrow progenitor cells in a similar way to other herpesviruses as CMV. However, ciHHV-6 represents an alternative form of viral persistence, occurring in a subgroup of individuals and characterized by very high viral loads in the blood ( $>1 \times 10^6$  HHV-6

copies/ml), in other body fluids or tissue samples. Growing evidence suggests that the integrated viral sequences are inherited through the germline and are therefore present in every nucleated cell. This phenomenon may confound the laboratory diagnosis of active HHV-6 infection because whole blood and spinal fluid from individuals with viral integration are persistently positive for HHV-6 DNA even in the absence of independent viral replication. Therefore, ciHHV-6 must be suspected every time whole blood HHV-6 DNA is detected at high load at the time of engraftment and persists at a high level after transplant, without any adverse influence on patients outcome.

Although the frequency of HHV-6 infection is quite different between allo-HSCT and ASCT, being higher after allo-HSCT,<sup>20</sup> the recognition of these cases may lead to the avoidance of unnecessary and potentially toxic antiviral treatments, particularly during the delicate phase of hematological recovery in the early post-transplant period. Currently, the screening for ciHHV-6 is not recommended, but we believe that the awareness of this phenomenon after ASCT is of interest for clinicians working in the field, since over 20,000 ASCTs are reported by the European Blood and Marrow Transplantation,<sup>21</sup> with an estimated number of 200-400 such new cases every year only in Europe.

**Acknowledgments.** We thank all the personnel of the Transplant Unit in the Oncology Department, at Humanitas Cancer Center, Rozzano, Italy, for patient's assistance, clinical care and follow-up.

## References:

1. [http://www.eortc.org/sites/default/files/ECIL3\\_CMV\\_HHV-6\\_Guidelines\\_Update\\_2009.pdf](http://www.eortc.org/sites/default/files/ECIL3_CMV_HHV-6_Guidelines_Update_2009.pdf)
2. Zerr DM. Human herpesvirus 6 and central nervous system disease in hematopoietic cell transplantation. *J Clin Virol.* 2006;37 Suppl 1:S52-56. [https://doi.org/10.1016/S1386-6532\(06\)70012-9](https://doi.org/10.1016/S1386-6532(06)70012-9)
3. Dulery R, Salleron J, Dewilde A, Rossignol J, Boyle EM, Gay J, de Berranger E, Coiteux V, Jouet JP, Duhamel A, Yakoub-Agha I. Early human herpesvirus type 6 reactivation after allogeneic stem cell transplantation: a large-scale clinical study. *Biol Blood Marrow Transplant.* 2012;18:1080-1089. <https://doi.org/10.1016/j.bbmt.2011.12.579> PMID:22212513
4. Sashihara J, Tanaka-Taya K, Tanaka S, Amo K, Miyagawa H, Hosoi G, Taniguchi T, Fukui T, Kasuga N, Aono T, Sako M, Hara J, Yamanishi K, Okada S. High incidence of human herpesvirus 6 infection with a high viral load in cord blood stem cell transplant recipients. *Blood.* 2002;100:2005-2511. PMID:12200359
5. Tormo N, Solano C, de la Cámara R, Garcia-Noblejas A, Carde-oso L, Clari MA, Nieto J, López J, Hernández-Boluda JC, Remigia MJ, Benet I, Navarro D. An assessment of the effect of human herpesvirus-6 replication on active cytomegalovirus infection after allogeneic stem cell transplantation. *Biol Blood Marrow Transplant.* 2010;16:653-661. <https://doi.org/10.1016/j.bbmt.2009.12.003> PMID:20005968
6. Fule Robles JD, Cheuk DK, Ha SY, Chiang AK, Chan GC. Human herpesvirus types 6 and 7 infection in pediatric hematopoietic stem cell transplant recipients. *Ann Transplant.* 2014;19:269-276. <https://doi.org/10.12659/AOT.889995> PMID:24881673
7. Tanaka-Taya K, Sashihara J, Kurahashi H, Amo K, Miyagawa H, Kondo K, Okada S, Yamanishi K. Human herpesvirus 6 (HHV-6) is transmitted from parent to child in an integrated form and characterization of cases with chromosomally integrated HHV-6 DNA. *J Med Virol.* 2004;73:465-473. <https://doi.org/10.1002/jmv.20113> PMID:15170644
8. Daibata M, Taguchi T, Nemoto Y, Taguchi H, Miyoshi I. Inheritance of chromosomally integrated human herpesvirus 6 DNA. *Blood.* 1999;94:1545-1549. PMID:10477678
9. Hubacek P, Virgili A, Ward KN, Pohlreich D, Keslova P, Goldova B, Markova M, Zajac M, Cinek O, Nacheva EP, Sedlacek P, Cetkovsky P. HHV-6 DNA throughout the tissues of two stem cell transplant patients with chromosomally integrated HHV-6 and fatal CMV pneumonitis. *Br J Haematol.* 2009;145:394-398. <https://doi.org/10.1111/j.1365-2141.2009.07622.x> PMID:19222466

10. Kamble RT, Clark DA, Leong HN, Heslop HE, Brenner MK, Carrum G. Transmission of integrated human herpesvirus-6 in allogeneic hematopoietic stem cell transplantation. *Bone Marrow Transplant.* 2007;40:563-566. <https://doi.org/10.1038/sj.bmt.1705780> PMID:17637686
11. Musso M, Scalone R, Marcacci G, Lanza F, Di Renzo N, Cascavilla N, Di Bartolomeo P, Crescimanno A, Perrone T, Pinto A. Fotemustine plus etoposide, cytarabine and melphalan (FEAM) as a new conditioning regimen for lymphoma patients undergoing auto-SCT: a multicenter feasibility study. *Bone Marrow Transplant.* 2010;45:1147-1153. <https://doi.org/10.1038/bmt.2009.318> PMID:19898504
12. De Pauw B, Walsh TJ, Donnelly JP, Stevens DA, Edwards JE, Calandra T, Pappas PG, Maertens J, Lortholary O, Kauffman CA, Denning DW, Patterson TF, Maschmeyer G, Bille J, Dismukes WE, Herbrecht R, Hope WW, Kibbler CC, Kullberg BJ, Marr KA, Mu-oz P, Odds FC, Perfect JR, Restrepo A, Ruhnke M, Segal BH, Sobel JD, Sorrell TC, Viscoli C, Wingard JR, Zaoutis T, Bennett JE; European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group; National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. Revised definitions of invasive fungal disease from the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. *Clin Infect Dis.* 2008;46:1813-1821. <https://doi.org/10.1086/588660> PMID:18462102 PMID:PMC2671227
13. Chapenko, Trociukas I, Donina S, Chistyakov M, Sultanova A, Gravelsina S, Lejniece S, Murovska M. Relationship between beta-herpesviruses reactivation and development of complications after autologous peripheral blood stem cell transplantation. *J Med Virol.* 2012;84:1953-1960. <https://doi.org/10.1002/jmv.23412> PMID:23080502
14. Piukovics K, Borbényi Z, Rajda C, Csomor A, Déak J, Terhes G. Monitoring human herpesvirus-6 in patients with autologous stem cell transplantation. *In Vivo* 2014;28:1113-1117. PMID:25398808
15. Colombier MA, Amorim S, Salmona M, Thieblemont C, Legoff J, Lafaurie M. HHV-6 reactivation as a cause of fever in autologous hematopoietic stem cell transplant recipients. *J Infect* 2017;75:155-159. <https://doi.org/10.1016/j.jinf.2017.05.011> PMID:28551368
16. Pellett PE, Ablashi DV, Ambros PF, Agut H, Caserta MT, Descamps V, Flamand L, Gautheret-Dejean A, Hall CB, Kamble RT, Kuehl U, Lassner D, Lautenschlager I, Loomis KS, Luppi M, Lusso P, Medveczky PG, Montoya JG, Mori Y, Ogata M, Pritchett JC, Rogez S, Seto E, Ward KN, Yoshikawa T, Razonable RR. Chromosomally integrated human herpesvirus 6: questions and answers. *Rev Med Virol* 2012;22:144-155. <https://doi.org/10.1002/rmv.715> PMID:22052666 PMID:PMC3498727
17. Kamble RT, Clark DA, Leong HN, Heslop HE, Brenner MK, Carrum G. Transmission of integrated human herpesvirus-6 in allogeneic hematopoietic stem cell transplantation. *Bone Marrow Transplant.* 2007;40:563-566. <https://doi.org/10.1038/sj.bmt.1705780> PMID:17637686
18. Clark DA, Nacheva EP, Leong HN, Brazma D, Li YT, Tsao EH, Buyck HC, Atkinson CE, Lawson HM, Potter MN, Griffiths PD. Transmission of integrated human herpesvirus 6 through stem cell transplantation: implications for laboratory diagnosis. *J Infect Dis.* 2006;193:912-916. <https://doi.org/10.1086/500838> PMID:16518751
19. Purev E, Winkler T, Danner RL, Fahle GA, Cook L, Zerr DM, Jerome KR, Childs RW. Engraftment of donor cells with germ-line integration of HHV6 mimics HHV6 reactivation following cord blood/haplo transplantation. *Blood.* 2014;124:1198-1199. <https://doi.org/10.1182/blood-2014-06-577684> PMID:25124787 PMID:PMC4133492
20. Quintela A, Escuret V, Roux S, Bonnafous P, Gilis L, Barraco F, Labussière-Wallet H, Duscassel-Leprêtre S, Nicolini FE, Thomas X, Chidiac C, Ferry T, Frobert E, Morisset S, Poitevin-Later F, Monneret G, Michallet M, Ader F; Lyon HEMINF Study Group. HHV-6 infection after allogeneic hematopoietic stem cell transplantation: from chromosomal integration to viral co-infections and T-cell reconstitution patterns. *J Infect* 2016 ;72 :214-222. <https://doi.org/10.1016/j.jinf.2015.09.039> PMID:26518057
21. Passweg JR, Baldomero H, Bader P, Bonini C, Cesaro S, Dreger P, Duarte RF, Dufour C, Kuball J, Farge-Bancel D, Gennery A, Kröger N, Lanza F, Nagler A, Sureda A, Mohty M. Hematopoietic stem cell transplantation in Europe 2014: more than 40000 transplants annually. *Bone Marrow Transplant.* 2016;51:786-792. <https://doi.org/10.1038/bmt.2016.20> PMID:26901709 PMID:PMC4895175