



## Original Article

### Liver Iron Content in Individuals With B-Thalassemia Trait and Hyperferritinemia: Role of Metabolic Alterations, *HFE* Genotypes, and Cirrhosis

Giulia Risca<sup>1,2</sup>, Raffaella Mariani<sup>3,4</sup>, Mara Botti<sup>3,4</sup>, Sara Pelucchi<sup>2</sup>, Stefania Galimberti<sup>1,5</sup> and Alberto Piperno<sup>4,6</sup>.

<sup>1</sup> Bicocca Bioinformatics Biostatistics and Bioimaging B4 Centre, University of Milano-Bicocca, Monza, Italy.

<sup>2</sup> Department of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy.

<sup>3</sup> Rare Disease Unit, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy.

<sup>4</sup> European Reference Network for Rare Haematological Diseases, EuroBloodNet.

<sup>5</sup> Biostatistics and Clinical Epidemiology, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy.

<sup>6</sup> Centro Ricerca Tettamanti, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italia.

**Competing interests:** The authors declare no competing interest.

## Supplementary Files

**Supplementary Table 1.** Classes of steatotic liver disease (SLD) according to LIC<sup>MRI</sup> grades.

Characteristics	LIC <sup>MRI</sup>				p-value
	Overall n=66	Grade 1 n=29 (43.9%)	Grade 2 n=24 (36.4%)	Grade 3 n=13 (19.7%)	
MASLD-MetALD	33 (50.0)	20 (69.0)	10 (41.7)	3 (23.1)	0.019
ALD	2 (3.0)	1 (3.4)	0	1 (7.7)	
Crypto SLD	2 (3.0)	1 (3.4)	1 (4.2)	0	
No SLD	29 (43.9)	7 (24.1)	13 (54.2)	9 (69.2)	

Legend: MASLD: metabolic-associated steatotic liver disease; MetALD: MASLD with moderate alcohol intake; ALD: alcohol-related steatotic liver disease; Crypto SLD: cryptogenic steatotic liver disease.